

Single production of excited spin-3/2 neutrinos at linear colliders

O. Çakır*

*Physics Department, Theory Division,
CERN, 1211 Geneva 23, Switzerland and
Ankara University, Faculty of Sciences,
Department of Physics, 06100, Tandoğan, Ankara, Turkey*

A. Ozansoy†

*Physics Department, CERN, 1211 Geneva 23, Switzerland and
Ankara University, Faculty of Sciences,
Department of Physics, 06100, Tandoğan, Ankara, Turkey*

Abstract

We study the potential of future high energy e^+e^- colliders to probe excited neutrino signals in different channels coming from the single production process via gauge interactions. We calculate the production cross section, decay widths and branching ratios of excited spin-3/2 neutrinos according to their effective currents and we compare them with that of the spin-1/2 ones. The signals and corresponding backgrounds are examined in detail to get accessible limits on the masses and couplings of excited spin-3/2 neutrinos.

*Electronic address: ocakir@mail.cern.ch

†Electronic address: aozansoy@mail.cern.ch

I. INTRODUCTION

Phenomenologically, an excited lepton can be considered as a heavy lepton sharing leptonic quantum number (flavour) with the corresponding Standard Model (SM) lepton. Excited leptons (l^*) are suggested by a composite model of leptons [1]. The lepton compositeness would be the most pronounced in the excess of high- p_T events in the forthcoming high energy experiments. If leptons are composites, they can be assigned to spin-1/2 bound states, containing three spin-1/2 or spin-1/2 and spin-0 subparticles. Bound states and/or excited states of spin-3/2 leptons are possible with three spin-1/2 [1] or spin-1/2 and spin-1 subparticles in the framework of compositeness. Composite leptons in the enlarged groups of standard theory would also imply spin-3/2 leptons [2, 3]. Further motivation for spin-3/2 particles comes from the supergravity (SUGRA) where spin-3/2 gravitino is the superpartner of the graviton [4]. Massive spin-3/2 excited neutrinos with the analogous heavy spin-1/2 excited ones can be produced at future high energy colliders through their effective interactions with the ordinary leptons. In the electron-positron collisions, pair production of excited spin-3/2 neutrinos can also be performed, but it is limited kinematically by $m^* < \sqrt{s}/2$, while single production directly can reach masses as high as $\sqrt{s}$. Furthermore, excited neutrino could contribute to the pair production of charged weak bosons through its exchange in the t -channel [5].

The mass limits for excited spin-1/2 neutrino from its single production search are $m^* > 190$ GeV and $m^* > 102.6$ GeV from their pair production [6] assuming $f = -f'$. Relatively small mass limits are obtained for $f = f'$, where f and f' are the scaling factors for the gauge couplings of $SU(2)$ and $U(1)$. In this case, the mass limits for the excited spin-1/2 neutrinos are: $m_{\nu_e^*} > 101.7$ GeV, $m_{\nu_\mu^*} > 101.8$ GeV and $m_{\nu_\tau^*} > 92.9$ GeV [6]. Recently, a search for the excited spin-1/2 neutrinos has been performed by H1 Collaboration [7] assuming $f = f'$ and $f/\Lambda = 1/m^*$, with an exclusion limit for the mass range of excited neutrino $m^* < 213$ GeV at 95% C.L. Excited spin-1/2 leptons was studied at hadron colliders in [8] and [9, 10] by taking into account possible backgrounds. These studies have shown that excited spin-1/2 lepton masses up to 2 TeV can be probed at the LHC. An analysis of the production and decay processes of single heavy spin-3/2 neutrinos was performed in [11, 12] in the frame of two phenomenological currents without taking into account possible backgrounds from the collisions.

In this study, we consider excited neutrino production in more detail (since excited spin-3/2 neutrinos are least studied compared to the excited spin-1/2 ones). We take into account the signals in different decay channels of excited neutrinos as well as the corresponding backgrounds at the International Linear Collider (ILC) [13] with $\sqrt{s} = 0.5$ TeV and Compact Linear Collider (CLIC) [14] with an optimal design energy of $\sqrt{s} = 3$ TeV. We present the missing transverse momentum distributions for single production of excited spin-3/2 and spin-1/2 neutrinos. The shape of these distributions would help to discriminate the signals from the background.

II. PHENOMENOLOGICAL CURRENTS

First, we consider the interaction between a spin-1/2 excited neutrino, gauge boson ($V = \gamma, Z, W^\pm$) and the SM lepton described by the effective current:

$$J_{1/2}^\mu = \frac{g_e}{2\Lambda} \bar{u}(k, 1/2) i\sigma^{\mu\nu} q_\nu (1 - \gamma_5) f_V u(p, 1/2) \quad (1)$$

where Λ is the scale of new physics responsible for the new interactions, k, p and q are the four-momentum of SM neutrino (electron), excited spin-1/2 neutrino and gauge boson γ, Z ($W^\pm$), respectively. g_e is the electromagnetic coupling constant, $g_e = \sqrt{4\pi\alpha}$. f_V is the electroweak coupling parameter corresponding to the vector boson V . In Eq. (1), $\sigma^{\mu\nu} = i(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu)/2$ with γ^μ being the Dirac matrices. For an excited spin-1/2 neutrino, three decay channels are possible: radiative decay $\nu^* \rightarrow \nu\gamma$, neutral weak decay $\nu^* \rightarrow \nu Z$, charged weak decay $\nu^* \rightarrow eW$. Neglecting the ordinary lepton masses we find decay widths as

$$\Gamma(l^{*(1/2)} \longrightarrow lV) = \frac{\alpha m^{*3}}{4\Lambda^2} f_V^2 \left(1 - \frac{m_V^2}{m^{*2}}\right)^2 \left(1 + \frac{m_V^2}{2m^{*2}}\right) \quad (2)$$

where $f_\gamma = (f - f')/2$; $f_Z = (f \cot \theta_W + f' \tan \theta_W)/2$; $f_W = f/\sqrt{2} \sin \theta_W$; m_V is the mass of the gauge boson. The parameters f and f' are determined by the composite dynamics, and they can be defined as q^2 -dependent form factors. In the literature, they are often taken as $f = f' = 1$ or $f = -f' = 1$ with $\Lambda = m^*$. The total decay width of the excited spin-1/2 neutrino is $\Gamma = 3.4(6.9)$ GeV for $m^* = 0.5(1)$ TeV at $f = f' = 1$ and $\Lambda = m^*$. The decay widths for excited spin-1/2 neutrino for the mass range $0.2 - 3$ TeV are given in Table I. The branching ratios of the excited spin-1/2 neutrinos into SM leptons and gauge bosons are given in Fig. 1. One may note that the electromagnetic interaction between

excited neutrino and ordinary neutrino vanishes for $f = f'$. As seen from Fig. 1, for $f = f' = 1$ charged current decays become dominant for higher mass values $m^* > 150$ GeV, while $f = -f' = 1$ the branching ratio is $\sim 60\%$ for eW channel. Their relative importance depends on the gauge boson mass and couplings. At $f = -f' = 1$ the branchings will be 60% for W channel, 12% for Z channel and 28% for γ channel at higher excited spin-1/2 neutrino masses ($m^* > 500$ GeV). Therefore the signature of $\nu^* \rightarrow eW$ is preferable in both cases for the investigation of excited neutrino in future linear collider experiments.




FIG. 1: The branching ratios (%) depending on the mass of excited spin-1/2 neutrino for $f = f' = 1$ (thin lines) and $f = -f' = 1$ (thick lines).

Second, the phenomenological currents for the interactions among spin-3/2 excited neutrino, gauge boson and SM lepton are given by

$$J_1^\mu = g_e u(k, 1/2) (c_{1V} - c_{1A} \gamma_5) u^\mu(p, 3/2) \quad (3)$$

$$J_2^\mu = \frac{g_e}{\Lambda} u(k, 1/2) q_\lambda \gamma^\mu (c_{2V} - c_{2A} \gamma_5) u^\lambda(p, 3/2) \quad (4)$$

$$J_3^\mu = \frac{g_e}{\Lambda^2} u(k, 1/2) q_\lambda i \sigma^{\mu\nu} q_\nu (c_{3V} - c_{3A} \gamma_5) u^\lambda(p, 3/2) \quad (5)$$

where $u^\mu(p, 3/2)$ represents the Rarita-Schwinger vector-spinor [15], c_{iV} and c_{iA} are the vector and axial vector couplings, the four momenta $q_\alpha = (p - k)_\alpha$ belongs to the vector boson. A spin-3/2 excited neutrino (ν^*) decays via two-body process according to the phenomenological currents Eq.(3-5). The radiative decay widths of excited spin-3/2 neutrinos for three

different currents are given as

$$\Gamma_1(\nu^{*(3/2)} \longrightarrow \nu\gamma) = \frac{\alpha}{4}(c_{1V}^2 + c_{1A}^2)m^* \quad (6)$$

$$\Gamma_2(\nu^{*(3/2)} \longrightarrow \nu\gamma) = \frac{\alpha}{24}(c_{2V}^2 + c_{2A}^2)m^*\left(\frac{m^*}{\Lambda}\right)^2 \quad (7)$$

$$\Gamma_3(\nu^{*(3/2)} \longrightarrow \nu\gamma) = \frac{\alpha}{48}(c_{3V}^2 + c_{3A}^2)m^*\left(\frac{m^*}{\Lambda}\right)^4 \quad (8)$$

and the weak decay widths of excited spin-3/2 neutrinos are given by

$$\Gamma_1(\nu^{*(3/2)} \longrightarrow lV) = \frac{\alpha}{48}(c_{1V}^2 + c_{1A}^2)m^*\frac{(1-\kappa)^2}{\kappa}(1+10\kappa+\kappa^2) \quad (9)$$

$$\Gamma_2(\nu^{*(3/2)} \longrightarrow lV) = \frac{\alpha}{48}(c_{2V}^2 + c_{2A}^2)m^*\left(\frac{m^*}{\Lambda}\right)^2\frac{(1-\kappa)^4}{\kappa}(1+2\kappa) \quad (10)$$

$$\Gamma_3(\nu^{*(3/2)} \longrightarrow lV) = \frac{\alpha}{48}(c_{3V}^2 + c_{3A}^2)m^*\left(\frac{m^*}{\Lambda}\right)^4(1-\kappa)^4(2+\kappa) \quad (11)$$

where $\kappa = (m_V/m^*)^2$ and m_V is the mass of the vector boson ($W^\pm$ or Z).

The relative importance of higher dimension operators is not essential when $\Lambda > m^*$. Taking $\Lambda = m^* = 0.5(1)$ TeV and $c_{iV} = c_{iA} = 0.5$ we find the total decay widths of the excited spin-3/2 neutrinos as $\Gamma_1 = 3.9(24.6)$ GeV, $\Gamma_2 = 2.7(22.2)$ GeV and $\Gamma_3 = 0.23(0.48)$ GeV for the currents J_1 , J_2 and J_3 , respectively. The decay widths of excited spin-3/2 neutrino with $c_{iV} = c_{iA} = 0.5$ for different mass values are shown in Table I. One can notice that even at $\Lambda = m^*$ and equal couplings, the values of the decay widths become different. Because, there are different radiative contributions for each currents and the κ term effects differently these channels. The corresponding branchings are given in Fig. 2. For equal couplings $c_{iV} = c_{iA} = 0.5$ and $\Lambda = m^*$ the branchings for the weak decays corresponding to the current J_1 and J_2 becomes dominant for $m^* \gtrsim 200$ GeV. The radiative and the weak decay channels with the same couplings have equal probability for the current J_3 when $m^* \gtrsim 500$ GeV.

III. CROSS SECTIONS

The spin-1/2 and spin-3/2 excited neutrinos can be produced singly at future e^+e^- colliders, namely ILC and CLIC. The Feynman diagrams contributing to the single production of excited neutrinos via the s -channel Z, γ exchange and t -channel W exchange are shown in Fig. 3.




FIG. 2: The branching ratios (%) of excited spin-3/2 neutrinos depending on their mass values. The plots (up to down) correspond to the currents J_1 , J_2 and J_3 with couplings $c_{iV} = c_{iA}$ and $\Lambda = m^*$.

TABLE I: Excited spin-3/2 neutrino decay widths depending on their masses for $c_{iV} = c_{iA} = 0.5$. The numbers in the second column show the results for excited spin-1/2 neutrinos with $f = -f' = 1$ ($f = f' = 1$) at $\Lambda = m^*$.

m^* (TeV)	$\Gamma_{J(1/2)}$ (GeV)	$\Gamma_{J_1(3/2)}$ (GeV)	$\Gamma_{J_2(3/2)}$ (GeV)	$\Gamma_{J_3(3/2)}$ (GeV)
0.2	1.15 (1.03)	0.54	0.14	0.06
0.3	1.93 (1.85)	1.22	0.55	0.12
0.4	2.67 (2.61)	2.29	1.36	0.18
0.5	3.39 (3.35)	3.89	2.71	0.23
0.75	5.18 (5.15)	11.12	9.31	0.36
1.0	6.95 (6.93)	24.62	22.20	0.48
1.5	10.43 (10.45)	78.89	75.24	0.78
2.0	13.97 (13.98)	183.48	178.61	0.97
2.5	17.49(17.47)	355.16	349.07	1.22
3.0	20.99 (20.98)	650.72	603.41	1.46




FIG. 3: Feynman diagrams for single excited spin-3/2 or spin-1/2 neutrino production in e^+e^- collisions.

One may note that pair production of excited neutrino in the t -channel W exchange is also possible with the currents given in the previous section. In the s -channel pair production of excited neutrinos of any type could also be possible if their interactions with photon and Z boson are defined. On the other hand, the single production of ν_μ^* and ν_τ^* with the s -channel

process is possible with and without the lepton flavour violating coupling (LFV).

Recently, spin-1/2 excited neutrino have been studied at the ILC energy $\sqrt{s} = 0.5$ TeV in [16]. The $\nu^* \rightarrow eW$ signal has been searched for an easy identification and accurate measurements in a linear collider environment. The results show that spin-1/2 excited neutrino can be probed up to the mass $m^* = 450$ GeV in the charged weak decay channel when the coupling parameters are taken as $f = f' = 0.1$ for $\Lambda = m^*$.

In order to calculate the cross sections for $e^+e^- \rightarrow \nu_e^* \bar{\nu}_e$ processes for three phenomenological spin-3/2 currents J_1, J_2 and J_3 , we integrate over the Mandelstam variable t (or $\cos \theta$) of the differential cross section

$$\frac{d\sigma^{(k)}}{dt} = \frac{g_e^2}{24m^{*2} \pi s^2} \sum_{i \leq j=1,3} \frac{T_{ij}^{(k)}}{P_{ij}^{(k)}} \quad (12)$$

where we use the expressions $T_{ij}^{(k)}$ and $P_{ij}^{(k)}$ as given in the Appendix. Total cross sections as a function of excited neutrino mass are presented in Figs. 4-6 at center of mass energy $\sqrt{s} = 0.5$ and $\sqrt{s} = 3$ TeV.




FIG. 4: Excited neutrino production cross section as a function of the mass at the ILC energy $\sqrt{s} = 0.5$ TeV (left panel) and CLIC energy $\sqrt{s} = 3$ TeV (right panel). Solid, dotted, dashed and dot-dashed lines denote spin-1/2 current $J(1/2)$ for $f = -f' = 1$ and the spin-3/2 with J_1, J_2, J_3 currents for $c_{iV}^\gamma = c_{iA}^\gamma = 0.5$, respectively.

One can see from Figs. 4 and 5, the excited spin-3/2 neutrinos with current J_3 (having only c_i^γ or c_i^Z couplings) has a cross section larger than the other two currents (J_1 and J_2) when its mass below 1.8 TeV and when they are produced at $\sqrt{s} = 3$ TeV. The cross section for the $\nu^*(J_3)$ via t -channel W -exchange becomes more visible below $m^* < 1.4$ TeV,




FIG. 5: Excited neutrino production cross section as a function of the mass at ILC with $\sqrt{s} = 0.5$ TeV (left panel) and CLIC with $\sqrt{s} = 3$ TeV (right panel). Solid, dotted, dashed and dot-dashed lines denote spin-1/2 and spin-3/2 currents $J(1/2)$, J_1 , J_2 , J_3 currents, respectively. Here, we take $f = f' = 1$ and $c_{iV}^Z = c_{iA}^Z = 0.5$ with $\Lambda = m^*$.




FIG. 6: Excited neutrino production cross section as a function of the mass at ILC (left panel) and CLIC (right panel). Solid, dotted, dashed and dot-dashed lines denote spin-1/2 and spin-3/2 currents for $f = -f' = 1$ and $c_{iV}^W = c_{iA}^W = 0.5$, respectively. Here we take $\Lambda = m^*$.

as shown in Fig. 6. Depending on the couplings c_{iV}, c_{iA} their (currents) relative importance (changes in the cross section for the interested mass range) become more pronounced compared to the excited spin-1/2 neutrinos.

In order to differentiate the spin-3/2 and spin-1/2 excited neutrino signals we plot the differential cross sections as a function of p_T in the Figs. 7-9. The distributions of missing

p_T for the single production of the spin-1/2 and spin-3/2 excited neutrino show different behaviours. Furthermore, the production of the excited spin-3/2 neutrino with J_3 interactions leads to different missing p_T distributions than the others (J_1 and J_2). Concerning the final state we consider three decay channels of signal $\nu^* \rightarrow \nu\gamma$, $\nu^* \rightarrow e^-W^+$ and $\nu^* \rightarrow \nu Z$ for excited neutrino. These lead to the final state with $\gamma + \cancel{p}_T$, $e^\pm + 2j + \cancel{p}_T$ and $\ell^+\ell^- + \cancel{p}_T$ as




FIG. 7: The missing p_T distribution taking into account the coupling $c_{iV}^\gamma = c_{iA}^\gamma = 0.5$ for spin-3/2 currents and $f = -f' = 1$ for spin-1/2 current with $\Lambda = m^*$ where $m^* = 350$ GeV for ILC and $m^* = 1$ TeV for CLIC.




FIG. 8: The missing p_T distribution taking into account the coupling $c_{iV}^Z = c_{iA}^Z = 0.5$ for spin-3/2 currents and $f = f' = 1$ for spin-1/2 currents with $\Lambda = m^*$ where we take $m^* = 350$ GeV for ILC and $m^* = 1$ TeV for CLIC.

shown in Table II. At the ILC energy of $\sqrt{s} = 500$ GeV, the shape of the missing transverse




FIG. 9: The missing p_T distribution taking into account the coupling $c_{iV}^W = c_{iA}^W = 0.5$ for the currents $J_i(3/2)$ and $f = -f' = 1$ for $J(1/2)$ with $\Lambda = m^*$ where we used $m^* = 350$ GeV for ILC and $m^* = 1$ TeV for CLIC.

momentum $\not{p}_T$ for the signal concerning $e^+e^- \rightarrow W^+e^-\bar{\nu}$ process behaves different from that of the corresponding background in Fig. 10. However, at higher center of mass energies the currents $J_1(3/2)$ and $J_2(3/2)$ show similar behaviour with the background especially at low $\not{p}_T$ region.

IV. ANALYSIS

In order to perceive the excited neutrino signals from the background we can put some cuts on the final state observable particles, and a cut on the missing transverse momentum. In general, by applying suitable cuts the signal can be more pronounced over the background. For the acceptance of interested events we apply the following initial cuts:

$$p_T^{e,\gamma} > 20 \text{ GeV} \quad (13)$$

$$|\eta_{e,\gamma}| < 2.5 \quad (14)$$

where p_T is the transverse momentum of final state detectable particle, η denotes for pseudorapidity. Since the neutrino can be identified as the missing transverse momentum, we can also apply the same cut on the $\not{p}_T$. After applying these cuts we find the total cross section for the SM background as 1.062 (2.049) pb, 0.329 (2.083) pb and 0.918 (0.463) pb at $\sqrt{s} = 0.5(3)$ TeV for the $e^+e^- \rightarrow \nu\bar{\nu}\gamma$, $e^+e^- \rightarrow \nu\bar{\nu}Z$ and $e^+e^- \rightarrow e^-\nu W^+$ processes,

TABLE II: Final states for single excited neutrino production

Decay mode	Production $e^+e^- \rightarrow \bar{\nu}\nu^*$
Radiative	$\bar{\nu}\nu\gamma$
Charged-current	$\bar{\nu}e^-2j$
Neutral-current	$\bar{\nu}\nu l^+l^-$

TABLE III: The cross section (in pb) for spin-3/2 signal for $c_{iV}^\gamma = c_{iA}^\gamma = 0.5$ and spin-1/2 signal for $f = -f' = 1$ at $\sqrt{s} = 0.5$ TeV. Here we take $\Lambda = m^*$

m^* (GeV)	J (1/2)	$J_1(3/2)$	$J_2(3/2)$	$J_3(3/2)$
200	5.36×10^0	1.85×10^{-1}	6.49×10^{-1}	2.60×10^0
250	3.21×10^0	1.27×10^{-1}	1.76×10^{-1}	4.90×10^{-1}
300	2.05×10^0	8.50×10^{-2}	4.70×10^{-2}	9.90×10^{-2}
350	1.43×10^0	5.10×10^{-2}	1.10×10^{-2}	1.80×10^{-2}
400	9.50×10^{-1}	2.50×10^{-2}	1.17×10^{-3}	2.30×10^{-3}
475	3.30×10^{-1}	1.80×10^{-3}	5.10×10^{-6}	5.40×10^{-6}

respectively. The program CalcHEP [17] is used to calculate the background cross sections with suitable cuts. In the analysis, we take into account one coupling c_{iV}, c_{iA} is kept free while the others are assumed to vanish. We present the signal cross sections in Tables III, IV, V, VI. In Tables VII and VIII we give the cross sections for the signal and background in the relevant mass intervals.

We examine the single production of excited spin-3/2 and spin-1/2 neutrinos in the decay channels $\nu\gamma$, νZ and $e^\pm W^\mp$. Concerning the final states we choose W boson decay hadronically and Z boson decay leptonically, because of the uncertainty from the neutrinos in the final state. The corresponding backgrounds can be studied by applying the cuts on the transverse momentum and pseudorapidities of final state leptons and jets. Furthermore, a way of extracting the excited neutrino signal is to impose a cut $|m_{ejj} - m^*| < 25$ GeV on the ejj invariant mass for the charged weak decay channels of ν^* . This cut can be relaxed for higher mass values of ν^* . For $m^* > 1.5$ TeV we apply an invariant mass cut $m_{ejj} > 1$ TeV for a clean detection of the signal.

TABLE IV: The signal cross sections (in pb) for $c_{iV}^\gamma = c_{iA}^\gamma = 0.5$ and $f = -f' = 1$ for excited neutrinos at $\sqrt{s} = 3$ TeV. Here we take $\Lambda = m^*$.

m^* (GeV)	$J(1/2)$	$J_1(3/2)$	$J_2(3/2)$	$J_3(3/2)$
250	4.97×10^0	6.90×10^{-2}	1.73×10^1	1.35×10^3
500	1.13×10^0	2.00×10^{-2}	1.00×10^0	2.01×10^1
1000	2.80×10^{-1}	6.70×10^{-3}	4.60×10^{-2}	2.50×10^{-1}
1500	1.29×10^{-1}	3.50×10^{-3}	4.90×10^{-3}	1.40×10^{-2}
2000	7.60×10^{-2}	1.70×10^{-3}	5.10×10^{-4}	9.10×10^{-4}
2500	5.00×10^{-2}	5.00×10^{-4}	2.10×10^{-5}	2.70×10^{-5}
2750	4.20×10^{-2}	1.30×10^{-4}	1.14×10^{-6}	1.29×10^{-6}

TABLE V: The signal cross section (in pb) for $c_{iV}^Z = c_{iA}^Z = 0.5$ and $f = f' = 1$ for the excited neutrinos at $\sqrt{s} = 0.5$ TeV. Here we take $\Lambda = m^*$.

m^* (GeV)	$J(1/2)$	$J_1(3/2)$	$J_2(3/2)$	$J_3(3/2)$
200	5.68×10^0	7.00×10^{-2}	2.40×10^{-1}	9.90×10^{-1}
250	3.82×10^0	4.80×10^{-2}	6.70×10^{-2}	1.90×10^{-1}
300	2.61×10^0	3.20×10^{-2}	1.80×10^{-2}	3.80×10^{-2}
350	1.84×10^0	2.00×10^{-2}	4.20×10^{-3}	6.90×10^{-3}
400	1.27×10^0	9.50×10^{-3}	6.40×10^{-4}	8.80×10^{-4}
475	4.40×10^{-1}	6.80×10^{-4}	1.92×10^{-6}	2.07×10^{-6}

For the analysis we define the statistical significance (SS) of the signal as

$$SS = \frac{\sigma_S}{\sqrt{\sigma_B}} \sqrt{\epsilon \cdot L}$$

where L_{int} is the integrated luminosity of the collider and ϵ is the efficiency to detect the signal in the chosen channel. In the $\nu^* \rightarrow \nu\gamma$ and $\nu^* \rightarrow \nu Z$ channel, we have used the cross section for the signal and background to calculate the significance. Requiring $SS > 3$ the attainable mass range covers up to $m^* \approx 1.3 - 1.5$ TeV (as seen from Fig. 11-13) for an excited spin-3/2 neutrino at CLIC with $\sqrt{s} = 3$ TeV. For the $\nu^* \rightarrow eW$ channel, the values of the significance are evaluated at each mass values (invariant mass intervals) and the results are presented in Fig. 13 the discovery reach for excited spin-3/2 and spin-1/2 neutrinos. A

TABLE VI: The cross sections (in pb) for $c_{iV}^Z = c_{iA}^Z = 0.5$ and $f = f' = 1$ for excited neutrinos at $\sqrt{s} = 3$ TeV, and $\Lambda = m^*$

m^* (GeV)	$J(1/2)$	$J_1(3/2)$	$J_2(3/2)$	$J_3(3/2)$
250	5.95×10^0	2.50×10^{-2}	6.15×10^0	4.80×10^2
500	1.50×10^0	7.00×10^{-3}	3.60×10^{-1}	7.16×10^0
1000	3.90×10^{-1}	2.40×10^{-3}	1.60×10^{-2}	9.00×10^{-2}
1500	1.79×10^{-1}	1.30×10^{-3}	1.70×10^{-3}	4.80×10^{-3}
2000	1.05×10^{-1}	6.10×10^{-4}	1.80×10^{-4}	3.20×10^{-4}
2500	7.00×10^{-2}	1.80×10^{-4}	7.54×10^{-6}	9.74×10^{-6}
2750	5.90×10^{-2}	4.80×10^{-5}	4.10×10^{-7}	4.60×10^{-7}

TABLE VII: The signal and background cross sections (in pb) after the cuts. The signal cross section is given for $\Lambda = m^*$ and $c_{iV}^W = c_{iA}^W = 0.5$ for the spin-3/2 and $f = -f' = 1$ for the spin-1/2 at $\sqrt{s} = 0.5$ TeV. We have used $\Delta m = 50$ GeV.

m^* (GeV)	$J(1/2)$	$J_1(3/2)$	$J_2(3/2)$	$J_3(3/2)$	$\sigma_B(\Delta m)$
200	8.82×10^0	2.30×10^0	3.61×10^0	7.70×10^0	9.36×10^{-2}
250	5.81×10^0	1.83×10^0	2.39×10^0	2.75×10^0	9.12×10^{-2}
300	4.02×10^0	1.52×10^0	1.76×10^0	1.25×10^0	9.65×10^{-2}
350	2.82×10^0	1.26×10^0	1.32×10^0	6.60×10^{-1}	1.21×10^{-1}
400	1.97×10^0	9.50×10^{-1}	9.10×10^{-1}	3.60×10^{-1}	1.69×10^{-1}
475	6.70×10^{-1}	2.80×10^{-1}	2.10×10^{-1}	9.50×10^{-2}	1.23×10^{-1}

smaller coupling $c_{iV}^W = c_{iA}^W = 0.05$ can be reached in the $\nu^* \rightarrow eW$ channel. At the ILC, the attainable mass ranges for spin-3/2 charged currents can be covered up to the center of mass energy.

V. CONCLUSION

We consider only the gauge interactions of excited spin-1/2 and spin-3/2 particles with the SM particles. Our analysis show that spin-3/2 excited neutrinos can be easily separated from the spin-1/2 ones by examining both the transverse momentum distributions of final

TABLE VIII: The signal and background cross sections (in pb) after the cuts. The cross section is given for $\Lambda = m^*$ and $c_{iV}^W = c_{iA}^W = 0.5$ for the spin-3/2 and $f = -f' = 1$ for the spin-1/2 signal at $\sqrt{s} = 3$ TeV.

m^* (GeV)	$J(1/2)$	$J_1(3/2)$	$J_2(3/2)$	$J_3(3/2)$	$\sigma_B(\Delta m)$
250	9.01×10^0	1.23×10^0	2.08×10^1	1.54×10^3	1.96×10^{-2}
500	2.35×10^0	3.70×10^{-1}	1.70×10^0	2.69×10^1	1.69×10^{-2}
1000	6.00×10^{-1}	1.60×10^{-1}	2.70×10^{-1}	6.50×10^{-1}	1.00×10^{-2}
1500	2.76×10^{-1}	1.20×10^{-1}	1.50×10^{-1}	1.00×10^{-1}	1.83×10^{-1}
2000	1.62×10^{-1}	1.10×10^{-1}	1.20×10^{-1}	3.50×10^{-2}	1.83×10^{-1}
2500	1.08×10^{-1}	1.00×10^{-1}	1.00×10^{-1}	1.70×10^{-2}	1.83×10^{-1}
2750	9.00×10^{-2}	9.70×10^{-2}	9.00×10^{-2}	1.30×10^{-2}	1.83×10^{-1}




FIG. 10: Missing p_T distributions of background processes $e^+e^- \rightarrow \gamma\nu\bar{\nu}$, $e^+e^- \rightarrow Z\nu\bar{\nu}$ and $e^+e^- \rightarrow W^+e^-\bar{\nu}$ at ILC (left) and CLIC (right) energies.

state visible and invisible particles when they produced singly at future linear colliders. If polarized e^+ and e^- beams are used the chiral structure of the couplings can be identified and more precise measurements can be performed at the ILC and CLIC environment. A previous work on excited spin-3/2 electrons [18] shows that they can be distinguished from the spin-1/2 ones by analyzing the angular distribution of final state observable particles.

Excited neutrinos can come in three families, ν_e^* , ν_μ^* and ν_τ^* , this study for ν_e^* can also be enlarged by applying similar analysis to the excited muon-neutrinos and tau-neutrinos at the future high energy linear colliders, and their signal distribution would be very different




FIG. 11: Statistical significance as a function of excited neutrino mass for different spin-3/2 currents with the couplings $c_{iV}^\gamma = c_{iA}^\gamma = 0.5$, and spin-1/2 current with $f = -f' = 1$ at $\sqrt{s} = 0.5$ TeV (first panel) and $\sqrt{s} = 3$ TeV (second panel) using an integrated luminosity of $L_{int} = 200 \text{ fb}^{-1}$ for ILC and $L_{int} = 400 \text{ fb}^{-1}$ for CLIC energies.




FIG. 12: Statistical significance as a function of excited neutrino mass for different spin-3/2 currents with the couplings $c_{iV}^Z = c_{iA}^Z = 0.5$, and spin-1/2 current with $f = f' = 1$ at $\sqrt{s} = 0.5$ TeV (first panel) and $\sqrt{s} = 3$ TeV (second panel) using an integrated luminosity of $L_{int} = 200 \text{ fb}^{-1}$ for ILC and $L_{int} = 400 \text{ fb}^{-1}$ for CLIC energies.

from that for the ν_{e^*} when they are produced in the t -channel diagrams with the LFV.




FIG. 13: Statistical significance as a function of excited neutrino mass for different spin-3/2 currents with the couplings $c_{iV}^W = c_{iA}^W = 0.05$, and spin-1/2 current with $f = -f' = 1$ at $\sqrt{s} = 0.5$ TeV (first panel) and $\sqrt{s} = 3$ TeV (second panel) using an integrated luminosity of $L_{int} = 200 \text{ fb}^{-1}$ for ILC and $L_{int} = 400 \text{ fb}^{-1}$ for CLIC energies.

A. Appendix

In equation (12) the operator terms $T_{ij}^{(1)}$ and propagators $P_{ij}^{(1)}$ for the spin-3/2 excited neutrino interaction current J_1 are given as

$$\begin{aligned}
T_{11}^{(1)} &= -g_e^2 (c_{1A}^\gamma{}^2 + c_{1V}^\gamma{}^2) (m^{*2} - s) (-t(s+t) + m^{*2}(2s+t)) \\
T_{12}^{(1)} &= [-g_e g_z (m_z^2 - s) (c_{1A}^\gamma (c_{1V}^Z c_A^f m^{*2} s (m^{*2} - s - 2t) + c_{1A}^Z c_V^f (st(s+t) + m^{*4}(2s+t) \\
&\quad - m^{*2}(2s^2 + 2st + t^2))) + c_{1V}^\gamma (c_{1A}^Z c_A^f m^{*2} s (m^{*2} - s - 2t) \\
&\quad + c_{1V}^Z c_V^f (st(s+t) + m^{*4}(2s+t) - m^{*2}(2s^2 + 2st + t^2))))]/2 \\
T_{13}^{(1)} &= [-g_e g_w (c_{1A}^W - c_{1V}^W) (c_{1A}^\gamma - c_{1V}^\gamma) ((m^{*4} + st)(s+t) - m^{*2}(s^2 + t^2))]/4\sqrt{2} \\
T_{22}^{(1)} &= [-g_z^2 (4c_{1A}^Z c_{1V}^Z c_A^f c_V^f m^{*2} s (m^{*2} - s - 2t) \\
&\quad + (c_{1A}^Z{}^2 + c_{1V}^Z{}^2) ((c_A^{f2} + c_V^{f2}) (m^{*2} - s) (-t(s+t) + m^{*2}(2s+t))))]/4 \\
T_{23}^{(1)} &= [-g_e g_w ((c_{1A}^W - c_{1V}^W) (c_{1A}^Z - c_{1V}^Z) (c_A^f + c_V^f) (m_z^2 - s) \\
&\quad ((m^{*4} + st)(s+t) - m^{*2}(s^2 + t^2)))]/8\sqrt{2} \\
T_{33}^{(1)} &= [-g_w^2 ((c_{1A}^W{}^2 + c_{1V}^W{}^2) (-m^{*2} + t) (s(s+t) - m^{*2}(s+2t))) \\
&\quad + 2c_{1A}^W c_{1V}^W m^{*2} (m^{*2} - 2s - t)t]/4
\end{aligned}$$

and

$$\begin{aligned}
P_{11}^{(1)} &= 1/s^2, P_{12}^{(1)} = 1/s(m_z^4 + s^2 + m_z^2(-2s + \Gamma_Z^2)), P_{13}^{(1)} = 1/s(m_w^2 - t), \\
P_{22}^{(1)} &= 1/(m_z^4 + s^2 + m_z^2(-2s + \Gamma_Z^2)), \\
P_{23}^{(1)} &= 1/((m_w^2 - t)(m_z^4 + s^2 + m_z^2(-2s + \Gamma_Z^2))), P_{33}^{(1)} = 1/(m_w^2 - t)^2
\end{aligned}$$

The $T_{ij}^{(2)}$ and $P_{ij}^{(2)}$ terms for the current J_2 are given by

$$\begin{aligned}
T_{11}^{(2)} &= -g_e^2 (c_{2A}^{\gamma^2} + c_{2V}^{\gamma^2}) (m^{*2} - s)^2 (-s^2 - 2st - 2t^2 + m^{*2} (s + 2t)) / 2 \\
T_{12}^{(2)} &= [g_e g_z (m^{*2} - s)^2 (-m_z^2 + s) (c_{2A}^{\gamma} (c_{2V}^Z c_A^f s (m^{*2} - s - 2t) \\
&\quad + c_{2A}^Z c_V^f (-s^2 - 2st - 2t^2 + m^{*2} (s + 2t))) \\
&\quad + c_{2V}^{\gamma} (c_{2A}^Z c_A^f s (m^{*2} - s - 2t) + c_{2V}^Z c_V^f (-s^2 - 2st - 2t^2 + m^{*2} (s + 2t))))] / 4 \\
T_{13}^{(2)} &= [-g_e g_w (c_{2A}^W + c_{2V}^W) (c_{2A}^{\gamma} + c_{2V}^{\gamma}) (m^{*2} - s - t) (s + t) (m^{*4} - st - m^{*2} (s + t))] / 4\sqrt{2} \\
T_{22}^{(2)} &= [-g_z^2 (m^{*2} - s)^2 (-4c_{2A}^Z c_{2V}^Z c_A^f c_V^f s (-m^{*2} + s + 2t) \\
&\quad + (c_{2A}^Z{}^2 + c_{2V}^Z{}^2) ((c_A^{f2} + c_V^{f2}) (-s^2 - 2st - 2t^2 + m^{*2} (s + 2t))))] / 8 \\
T_{23}^{(2)} &= [-g_z g_w ((c_{2A}^W + c_{2V}^W) (c_{2A}^Z + c_{2V}^Z) (c_A^f + c_V^f) (m_z^2 - s) (s + t) \\
&\quad (-m^{*2} + s + t) (s + t) (-m^{*4} + st + m^{*2} (s + t)))] / 8\sqrt{2} \\
T_{33}^{(2)} &= [-g_w^2 ((-m^{*2} + t)^2 (2c_{2A}^W c_{2V}^W (m^{*2} - 2s - t)t \\
&\quad + (c_{2A}^W{}^2 + c_{2V}^W{}^2) (-2s^2 - 2st - t^2 + m^{*2} (2s + t))))] / 4
\end{aligned}$$

and

$$\begin{aligned}
P_{11}^{(2)} &= 1/s^2 \Lambda^2, P_{12}^{(2)} = 1/s (m_z^4 + s^2 + m_z^2 (-2s + \Gamma_Z^2)) \Lambda^2, P_{13}^{(2)} = 1/s (m_w^2 - t) \Lambda^2, \\
P_{22}^{(2)} &= 1/(m_z^4 + s^2 + m_z^2 (-2s + \Gamma_Z^2)) \Lambda^2, \\
P_{23}^{(2)} &= 1/((m_w^2 - t) (m_z^4 + s^2 + m_z^2 (-2s + \Gamma_Z^2))) \Lambda^2, P_{33}^{(2)} = 1/(m_w^2 - t)^2 \Lambda^2
\end{aligned}$$

The $T_{ij}^{(3)}$ and $P_{ij}^{(3)}$ terms for the current J_3 are given by

$$\begin{aligned}
T_{11}^{(3)} &= -g_e^2 (c_{3A}^{\gamma^2} + c_{3V}^{\gamma^2}) (m^{*2} - s)^2 (m^{*4} + 2t(s+t) - m^{*2}(s+2t)) / 2 \\
T_{12}^{(3)} &= [-g_e g_z (m^{*2} - s)^2 (-m_z^2 + s) (c_{3A}^{\gamma} (c_{3V}^Z c_A^f m^{*2} (m^{*2} - s - 2t) - \\
&\quad c_{3A}^Z c_V^f (m^{*4} + 2t(s+t) - m^{*2}(s+2t))) + c_{3V}^{\gamma} (c_{3A}^Z c_A^f m^{*2} (m^{*2} - s - 2t) - \\
&\quad c_{3V}^Z c_V^f (m^{*4} + 2t(s+t) - m^{*2}(s+2t))))] / 4 \\
T_{13}^{(3)} &= [-g_e g_w (c_{3A}^W - c_{3V}^W) (c_{3A}^{\gamma} - c_{3V}^{\gamma}) (m^{*2} - s - t) t (m^{*2}(s-t) + st + m^{*2}(-s+t))] / 4\sqrt{2} \\
T_{22}^{(3)} &= [-g_z^2 (m^{*2} - s)^2 s (4c_{3A}^Z c_{3V}^Z c_A^f c_V^f m^{*2} (-m^{*2} + s + 2t) \\
&\quad + (c_{3A}^Z{}^2 + c_{3V}^Z{}^2) ((c_A^f{}^2 + c_V^f{}^2) (m^{*4} + 2t(s+t) - m^{*2}(s+2t)))] / 8 \\
T_{23}^{(3)} &= [g_z g_w s ((c_{3A}^W - c_{3V}^W) (c_{3A}^Z - c_{3V}^Z) (c_A^f + c_V^f) (m_z^2 - s) t \\
&\quad (-m^{*2} + s + t) (-m^{*2}(s-t) + st + m^{*2}(-s+t)))] / 8\sqrt{2} \\
T_{33}^{(3)} &= [-g_w^2 ((-m^{*2} + t)^2 t (2c_{3A}^W c_{3V}^W m^{*2} (-m^{*2} + 2s + t) \\
&\quad + ((c_{3A}^W{}^2 + c_{3V}^W{}^2) (m^{*4} + 2s(s+t) - m^{*2}(2s+t))))] / 4
\end{aligned}$$

and

$$\begin{aligned}
P_{11}^{(3)} &= 1/s\Lambda^4, P_{12}^{(3)} = 1/(m_z^4 + s^2 + m_z^2(-2s + \Gamma_Z^2))\Lambda^4, P_{13}^{(3)} = 1/(t - m_w^2)\Lambda^4, \\
P_{22}^{(3)} &= 1/(m_z^4 + s^2 + m_z^2(-2s + \Gamma_Z^2))\Lambda^4, \\
P_{23}^{(3)} &= 1/((t - m_w^2)(m_z^4 + s^2 + m_z^2(-2s + \Gamma_Z^2)))\Lambda^4, P_{33}^{(3)} = 1/(t - m_w^2)^2\Lambda^4
\end{aligned}$$

ACKNOWLEDGMENTS

We thank A. Belyaev and A. Pukhov for discussions on the CalcHEP implementation. OC acknowledges the support from the CERN TH Division. AO would like to thank to the members of the Nuclear Theory Group of University of Wisconsin for their hospitality. AO acknowledges the support through the Scientific and Technical Research Council (TUBITAK) BIDEB-2214. This work was partially supported by the Turkish State Planning Organisation (DPT) under grant No DPT-2006K-120470.

-
- [1] H. Terazawa, Y. Chikashige, and K. Akama, Phys. Rev. D **15**, 480 (1977); Y. Ne'eman, Phys. Lett. **82B** 69 (1969); H. Terazawa, M. Yasue, K. Akama, and M. Hayashi, Phys. Lett. **112B**,

- 387 (1982)
- [2] J. Leite Lopes, J. A. Martins Simoes, and D. Spehler, Phys. Lett. **94B**, 367 (1980); Phys. Rev. D **23**, 797 (1981); Phys. Rev. D **25**, 1854 (1982)
 - [3] Y. Tosa and R. E. Marshak, Phys. Rev. D **32** 774 (1985)
 - [4] D. Z. Freedman, P. van Nieuwenhuizen and S. Ferrara, Phys. Rev. D **13** (1976) 3214
 - [5] R. Walsh and A.J. Ramalho, Phys. Rev. D **67**, 097702 (2003).
 - [6] P. Achard et. al., Phys. Lett. B **568** 23-24 (2003); P. Achard et. al., Phys. Lett. B **531** 39-51 (2003).
 - [7] F.D. Aaron et. al, arXiv:hep-ex/08021858 (2008)
 - [8] U. Baur, M. Spira and P. M. Zerwas, Phys. Rev. D **42**, 815 (1990).
 - [9] O.J.P. Eboli and S.M. Lietti, Phys. Rev. D **65**, 075003 (2002).
 - [10] O. Cakir et al., Eur. Phys. J. C **32**, s02, 1 (2003).
 - [11] S. R. Choudhury, R. G. Ellis, G. C. Joshi, Phys. Rev. D **31**, 2390 (1985).
 - [12] F. M. L. Almeida et al., Phys. Rev. D **53**, 7 (1996)
 - [13] G. A. Loew, Report from the ILC technical review committee, Slac-pub-10024 (2003); A comprehensive information about the fututre linear colliders can be found at the URL:<http://www.linearcollider.org>
 - [14] R. W. Assmann et al.; The CLIC Study Team, A 3 TeV e^+e^- Linear Collider Based on CLIC Technology, CERN-2000-008, Geneva 2000; E.Accomando *et al.* [CLIC Physics Working Group], Physics at the CLIC Multi-TeV Linear Collider, arXiv:hep-ph/0412251.
 - [15] W. Rarita, J. Schwinger, Phys. Rev. **60**, 61 (1941).
 - [16] O.Cakir, A.Yilmaz and S. Sultansoy, Phys. Rev. D **70**, 075011 (2004); O.Cakir, I. Turk Cakir and Z. Kirca, Phys. Rev. D **70**, 075017 (2004).
 - [17] A. Pukhov *et al.*, arXiv: hep-ph/9908288; A. Pukhov, arXiv: hep-ph/0412191.
 - [18] O.Cakir and A.Ozansoy, Phys. Rev. D **77**, 035002 (2008).